

Diagnostic Services - Mouse



Deutscher Platz 5e
04103 Leipzig
Tel. 0341 39 29 86 90

info@gvg-diagnostics.de
www.gvg-diagnostics.de

Health monitoring of mouse colonies can be conducted through a combination of pathology, serology, microbiology, parasitology and PCR. In quarantine situations, PCR panels can be paired with serological tests to provide deeper insights into previous, inactive infections when importing animals.

We provide different health monitoring programs for quarterly or yearly monitoring according FELASA recommendations and more. Also, any kind of custom monitoring is possible.

Different sample types can be used for the different diagnostic procedures:

S - Serum
BS - Blot spot
T- Tissue
F- Feces

DS - Dry swab
MS - Microbiology swab (medium swab)
DSS - Deep skin scrape
FS - Fur swab

VIRUSES

| Serology: ELISA | Sample type | Virology S | Virology L | Virology XL | Virology XXL |
|-----------------------------------------------------|-------------|------------|------------|-------------|--------------|
| Mouse Hepatitis Virus (MHV) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Mouse rotavirus (EDIM) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Murine norovirus (MNV) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Minute virus of mice (MVM) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Mouse parvovirus (MPV) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Theiler's murine encephalomyelitis virus (TMEV) | S, BS | ✓ | ✓ | ✓ | ✓ |
| Lymphocytic choriomeningitis virus (LCV) | S, BS | | ✓ | ✓ | ✓ |
| Murine Adenovirus type 1/ type 2 (MAD-1/2 (FL/K87)) | S, BS | | ✓ | ✓ | ✓ |
| Ectromelia Virus (ECT) | S, BS | | ✓ | ✓ | ✓ |
| Murine Pneumonia Virus (PVM) | S, BS | | ✓ | ✓ | ✓ |
| Mammalian orthoreovirus type 3 (REO-3) | S, BS | | ✓ | ✓ | ✓ |
| Sendai Virus (SEN) | S, BS | | ✓ | ✓ | ✓ |
| Mouse cytomegalovirus (CMV) | S, BS | | | ✓ | ✓ |
| Hantavirus | S, BS | | | | ✓ |
| Maus K Virus | S, BS | | | | ✓ |
| Polyomavirus Mouse | S, BS | | | | |
| Murine Herpesvirus 3 (MTV) | S, BS | | | | |
| Lactate dehydrogenase elevating virus (LDV) | S, BS | | | | |
| Mouse kidney parvovirus (MKPV) | S, BS | | | | |
| Astrovirus | S, BS | | | | |

| PCR | Sample type | Virology S | Virology L | Virology XL | Virology XXL |
|-------------------------------------------------|-------------|------------|------------|-------------|--------------|
| Mouse rotavirus (EDIM) | F, DS | | | | |
| Mouse Hepatitis Virus (MHV) | F, DS, T | | | | |
| Murine norovirus (MNV) | F, DS | | | | |
| Theiler's murine encephalomyelitis virus (TMEV) | F, DS | | | | |
| Astrovirus | F, T | | | | |
| Lactate dehydrogenase elevating virus (LDV) | F, DS | | | | |
| Lymphocytic choriomeningitis virus (LCV) | DS | | | | |
| Mouse mammary tumor virus | T | | | | |
| Pneumonia virus of mice (PVM) | F, DS | | | | |
| Mammalian orthoreovirus type 3 (REO-3) | F, DS | | | | |
| Orthohantavirus hantanense | F, DS | | | | |
| Orthohantavirus seoulense | F, DS | | | | |
| Orthohantavirus sinnombreense | F, DS | | | | |
| Prospect Hill virus | F, DS | | | | |
| Sendai virus (SEN) | T, DS | | | | |
| Mouse parvovirus (MPV) | F, T, DS | | | | |
| Minute virus of mice (MVM) | F, T, DS | | | | |
| Ectromelia virus (ECT) | F, T | | | | |
| Mouse cytomegalovirus (CMV) | DS | | | | |
| Murine Adenovirus type 1 (MAD-1) | F, DS | | | | |
| Murine Adenovirus type 2 (MAD-2) | F, DS | | | | |
| Mouse kidney parvovirus (MKPV) | F, T | | | | |
| Mouse Thymic Virus (MTV) | T, DS | | | | |
| Murine pneumotropic virus (K Virus) | F, DS | | | | |
| Murine polyomavirus | F, DS | | | | |

Diagnostic Services - Mouse



Deutscher Platz 5e
04103 Leipzig
Tel. 0341 39 29 86 90

info@gvg-diagnostics.de
www.gvg-diagnostics.de

Health monitoring of mouse colonies can be conducted through a combination of pathology, serology, microbiology, parasitology and PCR. In quarantine situations, PCR panels can be paired with serological tests to provide deeper insights into previous, inactive infections when importing animals.

We provide different health monitoring programs for quarterly or yearly monitoring according FELASA recommendations and more. Also, any kind of custom monitoring is possible.

Different sample types can be used for the different diagnostic procedures:

S - Serum
BS - Blot spot
T- Tissue
F- Feces

DS - Dry swab
MS - Microbiology swab (medium swab)
DSS - Deep skin scrape
FS - Fur swab

BACTERIA / FUNGI

| Culture / MALDI-TOF | Sample type | Microbiology S | Microbiology L | MicrobiologyXL | Microbiology XXL |
|------------------------------------|-------------|----------------|----------------|----------------|------------------|
| Microbiology (MB) | T, MS | ✓ | ✓ | ✓ | ✓ |
| Citrobacter rodentium | T, MS | ✓ | ✓ | ✓ | ✓ |
| Corynebacterium kutscheri | T, MS | ✓ | ✓ | ✓ | ✓ |
| Klebsiella oxytoca | T, MS | ✓ | ✓ | ✓ | ✓ |
| Klebsiella pneumoniae | T, MS | ✓ | ✓ | ✓ | ✓ |
| Pasteurellaceae | T, MS | ✓ | ✓ | ✓ | ✓ |
| Pseudomonas aeruginosa | T, MS | ✓ | ✓ | ✓ | ✓ |
| Salmonella spp. | T, MS | ✓ | ✓ | ✓ | ✓ |
| Staphylococcus aureus | T, MS | ✓ | ✓ | ✓ | ✓ |
| Streptococci, β-haem. (A, B, C, G) | T, MS | ✓ | ✓ | ✓ | ✓ |
| Streptococcus pneumoniae | T, MS | ✓ | ✓ | ✓ | ✓ |

| Culture / MALDI-TOF | Sample type | Microbiology S | Microbiology L | MicrobiologyXL | Microbiology XXL |
|---------------------|-------------|----------------|----------------|----------------|------------------|
| Swab, aerobic | MS | | | | |
| Swab, anaerobic | MS | | | | |
| Mycology | MS | | | | |

| ELISA | Sample type | Microbiology S | Microbiology L | MicrobiologyXL | Microbiology XXL |
|--------------------------------|-------------|----------------|----------------|----------------|------------------|
| Rodentibacter pneumotropicus | S, BS | | | | |
| Rodentibacter heyllii | S, BS | | | | |
| Pasteurella multocida | S, BS | | | | |
| Mannheimia haemolytica | S, BS | | | | |
| Streptobacillus moniliformis | S, BS | | | | |
| Clostridium piliforme | S, BS | | ✓ | ✓ | ✓ |
| Mycoplasma pulmonis | S, BS | | ✓ | ✓ | ✓ |
| Mycoplasma arthritis | S, BS | | | | ✓ |
| Filibacterium rodentium (CARB) | S, BS | | | | ✓ |

| PCR | Sample type | Microbiology S | Microbiology L | MicrobiologyXL | Microbiology XXL |
|--------------------------------|-------------|----------------|----------------|----------------|------------------|
| Citrobacter rodentium | F, DS | | ✓ | ✓ | ✓ |
| Clostridium piliforme | F, DS | | | | |
| Corynebacterium bovis | F, DS | | | | |
| Corynebacterium kutscheri | F, DS | | | | |
| Helicobacter spp. | | | | | |
| Helicobacter bilis | F, DS | ✓ | ✓ | ✓ | ✓ |
| Helicobacter hepaticus | | | | | |
| Helicobacter typhlonius | | | | | |
| Helicobacter rodentium | F, DS | | | | |
| Klebsiella oxytoca | F, DS | | | | |
| Klebsiella pneumoniae | F, DS | | | | |
| Mycoplasma pulmonis | DS | | | | |
| Pseudomonas aeruginosa | F, DS | | | | |
| Proteus mirabilis | F, DS | | | | |
| Rodentibacter heyllii | F, DS | | | | |
| Rodentibacter pneumotropicus | F, DS | | | | |
| Staphylococcus aureus | F, DS | | | | |
| Salmonella spp. | F, DS | | | | |
| Streptobacillus moniliformis | DS | | ✓ | ✓ | ✓ |
| Streptococcus agalactiae (B) | DS | | | | |
| Streptococcus canis (G) | DS | | | | |
| Streptococcus equi (C) | DS | | | | |
| Streptococcus pyogenes (A) | DS | | | | |
| Streptococcus pneumoniae | F, DS | | | | |
| Bordetella bronchiseptica | DS | | | | |
| Bordetella hinzii | DS | | | | |
| Bordetella pseudohinzii | DS | | | | |
| Filibacterium rodentium (CARB) | DS | | | | |
| Pneumocystis carinii | T | | | | ✓ |

Diagnostic Services - Mouse



Deutscher Platz 5e
04103 Leipzig
Tel. 0341 39 29 86 90

info@gvg-diagnostics.de
www.gvg-diagnostics.de

Health monitoring of mouse colonies can be conducted through a combination of pathology, serology, microbiology, parasitology and PCR. In quarantine situations, PCR panels can be paired with serological tests to provide deeper insights into previous, inactive infections when importing animals.

We provide different health monitoring programs for quarterly or yearly monitoring according FELASA recommendations and more. Also, any kind of custom monitoring is possible.

Different sample types can be used for the different diagnostic procedures:

S - Serum
BS - Blot spot
T- Tissue
F- Feces

DS - Dry swab
MS - Microbiology swab (medium swab)
DSS - Deep skin scrape
FS - Fur swab

PARASITES

| Microscopy, Flotation | Sample type | Parasitology S | Parasitology L | Parasitology XL | Parasitology XXL |
|-----------------------|-------------|----------------|----------------|-----------------|------------------|
| Endoparasites (ENP) | | | | | |
| Helminths | F | ✓ | ✓ | ✓ | ✓ |
| Protozoa | F | ✓ | ✓ | ✓ | ✓ |
| Ectoparasites (ECP) | FS | ✓ | ✓ | ✓ | ✓ |
| Scabies mites | DSS | | | | ✓ |

| ELISA | Sample type | Parasitology S | Parasitology L | Parasitology XL | Parasitology XXL |
|--------------------------|-------------|----------------|----------------|-----------------|------------------|
| Cryptosporidium muris | F | | | | ✓ |
| Encephalitozoon cuniculi | F | | | | ✓ |

| PCR | Sample type | Parasitology S | Parasitology L | Parasitology XL | Parasitology XXL |
|------------------------|-------------|----------------|----------------|-----------------|------------------|
| Aspicularis tetraptera | F | | | | |
| Eimeria spp. | F | | | | |
| Entamoeba muris | F | | | | |
| Cryptosporidium muris | F | | | | |
| Chilomastix spp. | F | | | | |
| Giardia muris | F | | | | ✓ |
| Hexamastix spp. | F | | | | |
| Spironucleus muris | F | | | | |
| Syphacia muris | F | | | | |
| Syphacia obvelata | F | | | | |
| Tritrichomonas muris | F | | | | |
| Myobia musculi | FS | | | | |
| Myocoptes musculinus | FS | | | | |
| Radfordia affinis | FS | | | | |

| Diagnostic service | Method | Pathology | Pathology | Pathology | Pathology |
|--------------------------------|------------|-----------|-----------|-----------|-----------|
| Postmortem examination | Necropsy | ✓ | ✓ | ✓ | ✓ |
| Histological exam. (per organ) | Microscopy | | | | |